










[Home](#)
[Quick](#)
[Advanced](#)
[Pat Num](#)
[Help](#)

10/642, 194.

Jump To

an/orchid and cephalosporin and sustained and relea

- 1 [6,903,211](#)  [Process for the preparation of 3-propenyl cephalosporin DMF solvate](#)
- 2 [6,803,461](#)  [Synthesis of ceftiofur intermediate](#)
- 3 [6,800,756](#)  [Method for the preparation of ceftiofur sodium and its intermediates](#)
- 4 [6,800,755](#)  [Process for the preparation of cefixime](#)
- 5 [6,713,625](#)  [Process for the preparation of cefditoren using the thioester of thiazolylacetic acid](#)
- 6 [6,610,845](#)  [Thioester derivatives of thiazolyl acetic acid and their use in the preparation of cephalosporin compounds](#)
- 7 [6,555,680](#)  [Method for the preparation of ceftiofur sodium](#)
- 8 [6,388,070](#)  [Thioester derivatives of thiazolyl acetic acid and their use in the preparation of cephalosporin compounds](#)
- 9 [6,384,215](#)  [Preparation of new intermediates and their use in manufacturing of cephalosporin compounds](#)

[Home](#)
[Quick](#)
[Advanced](#)
[Pat Num](#)
[Help](#)